

Biological and molecular effects of radiation

Chronic Radiation Effects

Healthy adults from High Level Natural Radiation Areas of Kerala








High Level Natural Radiation Area
 ≤ 1.0 mGy to >45 mGy/y






Normal Level Natural Radiation Area
 ≤ 1.5 mGy/y

Acute Radiation Effects

Random healthy adults from Mumbai



Comparative Proteomic Analysis: 2D PAGE/DIGE & LC-MS/MS



Mechanism of Radio-adaptive response



Primed cells show:

- Lower DNA damage & better repair
- Lower levels of ROS
- Higher activity and gene expression of antioxidant enzymes
- Increased binding of transcription factors Nrf2 and NFκB
- Enrichment of proteins involved in transcriptional, ubiquitination, signaling and chromatin remodelling.

HLNRA individuals show


- Evidence of radio-adaptive response.
- Higher expression of DNA repair & pro-survival proteins

Understanding mechanisms of radio-resistance in cancer cells




- A549 (Human lung carcinoma) cells were found to be relatively more radio-resistant with 10Gy dose delivered as a **fractionated** regimen.

y-H2AX, 15min y-H2AX, 4h





p-TP53 p-TP53 + DAPI



Radio-resistant cells show:

- Efficient DNA repair
- Upregulation of DNA repair pathway-associated genes (DNA-PK, ATM, Rad52, MLH1 and BRCA1).
- Translocation of phospho-p53 into the nucleus of A549 cells.




Silencing of DNA repair gene Rad 52 makes A549 cells radio-sensitivet


Radiation Signaling Group

Biological effectiveness of Low and High LET radiation


Proton irradiation



4 Gy of Gamma equitoxic to Proton 2 Gy





Transcriptional response of proton and gamma irradiated cells



Equitoxic dose of proton suppresses EMT and cancer stem cell like cells phenotype




Carbon - beam irradiation



Carbon-beam irradiated cells showed:

- 1.2 times higher γH2Ax foci than gamma.
- Qualitative and quantitative differences in Radiation induced foci (RIF) of ATM, ATR, BRCA1 compared to gamma.
- Early apoptosis







Oxygen - beam irradiation



Oxygen-irradiated cells showed:


- An RBE of 3 at 20% survival
- Differences in DNA damage response with equitoxic doses of low LET radiation

Radiation induced bystander signaling




Bystander cells showed:

- Up-regulated expression of NF-κB, iNOS, p53 and p21 genes
- Increased DNA damage, apoptosis and NO
- Bystander signaling between macrophages and EL-4 cells
- Reduced bystander response after L-NAME treatment
- Partially reduced bystander response after cPTIO treatment




(A) Control

(D) Medium from irradiated U937



(B) Cross Bystander

(C) Medium from stimulated U937





- Existence of bystander signaling between different cell types (U937 and A549 cells)
- Suppression of ATM with siRNA completely inhibits bystander effect in similar cell types

Radiation Signaling Group

Interaction of cold atmospheric pressure plasma with cancer cells

In collaboration with Institute for Plasma Research (IPR), Gandhinagar

Single Jet System




Plasma exposure reduces A549 cell viability in a dose-dependent manner

Multi-jet System




Cell Model

Human lung adenocarcinoma cell line (A549)



DNA damage analysis with alkaline comet assay



Plasma exposure leads to a significant increase in DNA damage in A549 cells